



## Medication Dosage Chart

Note: *Aspirin should not be given to any child*

<b>Acetaminophen (Tylenol)</b>	<b>Tylenol should be given every 4-6 hours. Do not dose more often than every 4 hours.</b>								
	1-3mo. 6-11 lbs	4-11mo. 12-17 lbs	12-23 mo. 18-23 lbs	2-3 yr. 24-35 lbs	4-5 yr. 36-47 lbs	6-8 yr. 48-59 lbs	9-11 yr 60-71 lbs	72-95 lbs	95 + lbs
Baby Drops 80mg/0.8ml	½ dropper 0.4 ml	1 dropper 0.8 ml	1 ½ dropper 1.2 ml	2 droppers 1.6 ml	2 ½ droppers 2.4 ml				
Elixir or Syrup 160mg/5ml	1.25 ml	2.5 ml	4ml	1 tsp	1.5 tsp	2 tsp	2.5 tsp	3 tsp	4 tsp
Chewable Tabs 80mg			1.5 tabs	2 tabs	3 tabs	4 tabs	5 tabs	6 tabs	8 tabs
Chewable Tabs 160 mg				1 tab	1.5 tab	2 tabs	2.5 tabs	3 tabs	4 tabs
Adult Tabs 325 mg				0.5 tab	0.75 tab	1 tab	1.25 tabs	1.5 tabs	2 tabs
<b>Ibuprofen (Motrin/Advil)</b>	<b>Do not give Ibuprofen to children under 6 months of age. Ibuprofen should be given every 6-8 hours. Do not give doses more often than every 6 hours.</b>								
		4-11mo. 12-17 lbs	12-23 mo. 18-23 lbs	2-3 yr. 24-35 lbs	4-5 yr. 36-47 lbs	6-8 yr. 48-59 lbs	9-11 yr 60-71 lbs	72-95 lbs	95 + lbs
Infant Drops 50mg/1.25ml		1.25 ml	1.875ml	2.5 ml	3.125 ml	3.725 ml			
Suspension 100mg/5ml		2.5 ml	4 ml	1tsp	1.5 tsp	2 tsp	2.75 tsp	3.25 tsp	4 tsp
Chewable Tabs 50mg		1 tab	1.5 tabs	2 tabs	3 tabs	4 tabs	5.5 tabs	6.5 tabs	8 tabs
Chewable Tabs 100mg		0.5 tab	1 tab	1 tab	1.5 tab	2 tabs	2.5 tabs	3 tabs	4 tabs
Caplets Jr. 100mg		0.5 tab	1 tab	1 tab	1.5 tab	2 tabs	2.5 tabs	3 tabs	4 tabs
Adult Tabs 200mg				0.5 tab	0.75 tab	1 tab	1 tab	1.5 tab	2 tab
<b>Diphehydramine (Benadryl)</b>	<b>Used for sedation and mild decongestant effect. May cause children to be more sleepy or more hyper than usual. Do not give doses more often than every 4 to 6 hours.</b>								
		6-11mo. 14-17 lbs	12-23 mo. 18-23 lbs	2-3 yr. 24-35 lbs	4-5 yr. 36-47 lbs	6-8 yr. 48-59 lbs	9-11 yr. 60-84 lbs	85-111 lbs	112 + lbs
Syrup 12.5mg/5ml		¾ tsp	1 tsp	1.2 tsp	1 ¾ tsp	2 tsp	2 tsp		
Chewables 12.5 mg					1 tab	2 tabs	3 tabs	3.5 tabs	
Adult Caps 25mg						1 cap	1 cap	2 cap	2.5 cap

\*1 tsp= 5cc or 5ml, ½ tsp=2.5cc or 2.5ml

**Note:** Consult with a physician or pharmacist before combining any medication. Some drugs may contain additional pain relievers.